

AP Review 17: Mixed Review 2

AB/BC Calc 2010 #2 Calc Ok

t (hours)	0	2	5	7	8
$E(t)$ (hundreds of entries)	0	4	13	21	23

A zoo sponsored a one-day contest to name a new baby elephant. Zoo visitors deposited entries in a special box between noon ($t = 0$) and 8 P.M. ($t = 8$). The number of entries in the box t hours after noon is modeled by a differentiable function E for $0 \leq t \leq 8$. Values of $E(t)$, in hundreds of entries, at various times t are shown in the table above.

(a) Use the data in the table to approximate the rate, in hundreds of entries per hour, at which entries were being deposited at time $t = 6$. Show the computations that lead to your answer.

(b) Use a trapezoidal sum with the four subintervals given by the table to approximate the value of $\frac{1}{8} \int_0^8 E(t) dt$.

Using correct units, explain the meaning of $\frac{1}{8} \int_0^8 E(t) dt$ in terms of the number of entries.

(c) At 8 P.M., volunteers began to process the entries. They processed the entries at a rate modeled by the function P , where $P(t) = t^3 - 30t^2 + 298t - 976$ hundreds of entries per hour for $8 \leq t \leq 12$. According to the model, how many entries had not yet been processed by midnight ($t = 12$)?

(d) According to the model from part (c), at what time were the entries being processed most quickly? Justify your answer.

BC Calc 2010 #5 No Calc (**BC ONLY**)

Consider the differential equation $\frac{dy}{dx} = 1 - y$. Let $y = f(x)$ be the particular solution to this differential equation with the initial condition $f(1) = 0$. For this particular solution, $f(x) < 1$ for all values of x .

- (a) Use Euler's method, starting at $x = 1$ with two steps of equal size, to approximate $f(0)$. Show the work that leads to your answer.
- (b) Find $\lim_{x \rightarrow 1} \frac{f(x)}{x^3 - 1}$. Show the work that leads to your answer.
- (c) Find the particular solution $y = f(x)$ to the differential equation $\frac{dy}{dx} = 1 - y$ with the initial condition $f(1) = 0$.

BC Calc 2003 Form B #6 No Calc (BC ONLY)

The function f has a Taylor series about $x = 2$ that converges to $f(x)$ for all x in the interval of convergence. The n th derivative of f at $x = 2$ is given by $f^{(n)}(2) = \frac{(n+1)!}{3^n}$ for $n \geq 1$, and $f(2) = 1$.

- (a) Write the first four terms and the general term of the Taylor series for f about $x = 2$.
- (b) Find the radius of convergence for the Taylor series for f about $x = 2$. Show the work that leads to your answer.
- (c) Let g be a function satisfying $g(2) = 3$ and $g'(x) = f(x)$ for all x . Write the first four terms and the general term of the Taylor series for g about $x = 2$.
- (d) Does the Taylor series for g as defined in part (c) converge at $x = -2$? Give a reason for your answer.

Homework

As usual, try the problems without notes, then score yourself using the videos and scoring guides. Make corrections in a different color and then fill out the score report form. Email me more your work with corrections. AB Test Takers need to on do the first TWO FRQs. BC Test Takers, need to do the LAST THREE FRQs. You may do the first, if you wish.

AB Calc 2007 #6 No Calc (**AB ONLY**)

Let f be the function defined by $f(x) = k\sqrt{x} - \ln x$ for $x > 0$, where k is a positive constant.

- (a) Find $f'(x)$ and $f''(x)$.
- (b) For what value of the constant k does f have a critical point at $x = 1$? For this value of k , determine whether f has a relative minimum, relative maximum, or neither at $x = 1$. Justify your answer.
- (c) For a certain value of the constant k , the graph of f has a point of inflection on the x -axis. Find this value of k .

AB/BC Calc 2003 #6 No Calc (AB/BC)

Let f be the function defined by

$$f(x) = \begin{cases} \sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ 5-x & \text{for } 3 < x \leq 5. \end{cases}$$

- (a) Is f continuous at $x = 3$? Explain why or why not.
(b) Find the average value of $f(x)$ on the closed interval $0 \leq x \leq 5$.
(c) Suppose the function g is defined by

$$g(x) = \begin{cases} k\sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ mx+2 & \text{for } 3 < x \leq 5, \end{cases}$$

where k and m are constants. If g is differentiable at $x = 3$, what are the values of k and m ?

BC Calc 2013 #6 No Calc (**BC ONLY**)

A function f has derivatives of all orders at $x = 0$. Let $P_n(x)$ denote the n th-degree Taylor polynomial for f about $x = 0$.

- (a) It is known that $f(0) = -4$ and that $P_1\left(\frac{1}{2}\right) = -3$. Show that $f'(0) = 2$.
- (b) It is known that $f''(0) = -\frac{2}{3}$ and $f'''(0) = \frac{1}{3}$. Find $P_3(x)$.
- (c) The function h has first derivative given by $h'(x) = f(2x)$. It is known that $h(0) = 7$. Find the third-degree Taylor polynomial for h about $x = 0$.

BC Calc 2008 #5 No Calc (**BC ONLY**)

The derivative of a function f is given by $f'(x) = (x - 3)e^x$ for $x > 0$, and $f(1) = 7$.

- The function f has a critical point at $x = 3$. At this point, does f have a relative minimum, a relative maximum, or neither? Justify your answer.
- On what intervals, if any, is the graph of f both decreasing and concave up? Explain your reasoning.
- Find the value of $f(3)$.